

Short communication

Rare sesquiterpenes from South African *Pteronia* species

A.M. Viljoen^{a,*}, G.P.P. Kamatou^a, Z.H. Coovadia^b, T. Özek^c, K.H.C. Başer^c

^a Department of Pharmaceutical Sciences, Faculty of Science, Tshwane University of Technology, Private Bag X680, Pretoria 0001, South Africa

^b Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, 7 York Road, Parktown 2193, South Africa

^c Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University, 26470-Eskişehir, Turkey

Received 19 June 2009; received in revised form 7 July 2009; accepted 30 July 2009

Abstract

The genus *Pteronia* consists of approximately 80 species which are widely distributed in southern Africa. The essential oils isolated from the aerial parts of eleven species, analyzed by GC-MS varied both qualitatively and quantitatively. In *Pteronia pallens*, *Pteronia empetrifolia* and *Pteronia flexicaulis* uncommon sesquiterpenes such as presilphiperfol-7-ene, 7- α -(H)-silphiperfol-5-ene, 7- β -(H)-silphiperfol-5-ene, α -campholene aldehyde, silphiperfol-5-ene, cameroonan-7- α -ol, silphiperfol-7- β -ol, presilphiperfolan-9- α -ol and presilphiperfolan-8-ol (a major compound in *P. pallens*) were identified. Cluster analysis based of the chemical composition of the oils revealed that individual plants of *Pteronia camphorata* collected in the same population had similar oil profiles with a high correlation coefficient ($S_{\text{corr}} \approx 0.98$). Similarly, the essential oil composition of *P. pallens* collected from two distinct localities also showed high levels of congruency ($S_{\text{corr}} \approx 0.99$).

© 2009 SAAB. Published by Elsevier B.V. All rights reserved.

Keywords: 7- α -(H)-silphiperfol-5-ene; 7- β -(H)-Silphiperfol-5-ene; Essential oil; Presilphiperfolan-8-ol; Presilphiperfolan-9- α -ol; Presilphiperfol-7-ene; *Pteronia*; GC-MS; Silphiperfol-5-ene; Silphiperfol-7- β -ol; α -Campholene aldehyde

1. Introduction

Pteronia (L.) L. (Asteraceae) is represented by approximately 80 species, which are primarily found in southern Africa. The species are most abundant in the Karoo-Namib region although 12 species occur in the Fynbos biome of South Africa (Goldblatt and Manning, 2000). Members of this genus are often the dominant taxa in the plant communities in which they are found.

The morphology of the genus is diverse with regards to habit, flowers and leaf shape. They are mainly perennial and woody shrubs ranging from 0.3 to 1.5 m in height. The leaf texture varies from smooth, lacking hair and bristles (glabrous) to hairy, with leaves either opposite or alternate (Shearing, 1997).

Two species; *Pteronia pallens* and *Pteronia paniculata* are unpalatable species that dominate the veld in large parts of the Little Karoo. Another well documented characteristic of *Pter-*

onia (e.g. *P. pallens*) is that it is highly toxic to ungulate herbivores, and may be lethal to cattle.

Aromatherapy (uses of volatiles from plants for improvement of mood or health) is characteristic of the San culture of South Africa. The name San was used to refer to one of the most preferred species *Pteronia ombromoides* (Van Wyk, 2008). Since ancient times, the Khoi-San have been using *Pteronia* species which have been amongst the first plants to be used for their anti-infective properties and have also been used in the treatment of stomach ailments (cramps) and nausea (Shearing, 1997). Many species have also shown anti-insecticidal properties (Shearing, 1997). Considering the extensive historic use of *Pteronia* in traditional medicine, it is ironic that the volatiles of this genus remain poorly investigated.

2. Material and methods

2.1. Plant studied

Eleven species (14 samples) were collected from various areas in the eastern and southern Cape of South Africa (Table 1). All

* Corresponding author. Tel.: +27 12 3826360; fax: +27 12 3826243.
E-mail address: viljoenam@tut.ac.za (A.M. Viljoen).

Table 1
List of species included in this study with voucher and locality information.

| Species | Voucher number | Location |
|-----------------------------|----------------|-------------------------------------|
| <i>P. adenocarpa</i> Harv. | Vlok 2822 | Frinsgeword |
| <i>P. camphorata</i> (L.)L. | Vlok 2804 | Montagu Pass near Herold |
| <i>P. elongate</i> Thunb. | Vlok 2819 | Prins Albert on Klaarstroom road |
| <i>P. empetrifolia</i> Dc. | Vlok 2818 | Prins Albert |
| <i>P. fasciculata</i> L.f. | Vlok 2814 | Bakenshoogte |
| <i>P. flexicaulis</i> L.f. | Vlok 2805 | Southern Hills near Daskop |
| <i>P. glauca</i> Thunb. | Vlok 2812 | Lategansvlei |
| <i>P. glomerata</i> L.f. | Vlok 2824 | West of Klaarstroom |
| <i>P. pallens</i> L.f. | Vlok 2815 | Volmoed |
| <i>P. pallens</i> L.f. | Vlok 2816 | Between Oudtshoorn and Volmoed |
| <i>P. paniculata</i> Thunb. | Vlok 2813 | Bakenshoogte |
| <i>P. viscosa</i> Thunb. | Vlok 2817 | Prins Albert |

plants were identified by J Vlok, who also assisted in preparing the voucher specimens which are housed in the Department of Pharmaceutical Sciences (Tshwane University of Technology, Pretoria). Three individual samples were collected in the same population for *Pteronia camphorata* and two samples of *Pteronia pallens* were collected at different localities to assess possible within and between population variation.

2.2. Isolation of the essential oils

The aerial part of each plant was air-dried at room temperature and the oil was isolated by hydro-distillation using a Clevenger apparatus together with 500 ml of water over 4 h. The resultant oil was stored at 4 °C until analysis was carried out.

2.3. Gas chromatography coupled to mass spectroscopy (GC-MS)

The analysis of the essential oil was performed using a Hewlett-Packard GCD system equipped with a HP-Innowax column (60 m × 0.25 mm Ø, with 0.25 µm film thickness). Temperature at the injection port was 250 °C. Column temperature was initially 60 °C for 10 min then gradually increased to 220 °C at a 4 °C/min. Temperatures were held at 220 °C for 10 min until it was finally raised to 240 °C at a rate of 1 °C/min. Helium was used as carrier gas at a flow rate of 0.7 mL/min. An electron ionization system with an ionization energy of 70 eV was used with a split ratio of 50:1. The mass range was between 35 and 425 m/z. Percentage compounds were calculated by the computer from total ion chromatograms. Component identifications were performed by comparing their mass spectra and retention indices using the Başer Library of Essential Oil Constituents.

The percentage composition of the essential oil samples was used to determine the similarity between samples by cluster analysis using the NTSYS software (Rohlf, 1992). Correlation was selected as a measure of similarity and the unweighted pair-group method with arithmetic average (UPGMA) was used for

cluster definition. The degree of correlation was evaluated according to Pestana and Gageiro (2000) where a very high correlation ranged between 0.90 and 1.00, high between 0.70 and 0.89, moderate between 0.40 and 0.69, low between 0.20 and 0.39 and very low if less than 0.20.

3. Results and discussion

The essential oil composition is given in Table 2. A total of 86 compounds representing 91.8% to 98.8% of the oil were identified. The following monoterpenes were present in all the oils; α-pinene (0.2–12.0%), β-pinene (0.7–32.1%), sabinene (2.0–33.6%), limonene (2.7–13.0%), p-cymene (0.5–21.1%), α-terpineol (0.1–4.8%) and terpinen-4-ol (0.8–9.2%). However, some major and minor compounds differ extensively within the genus *Pteronia* (Table 2). This confirms findings of Zedro et al. (1990) that the diterpenes and other compounds commonly present in *Pteronia* species are not uniform although there are some similarities.

In *P. pallens*, *Pteronia empetrifolia* and *Pteronia flexicaulis*, uncommon compounds characterise the essential oil; presilphiperfol-7-ene, 7-α-(H)-silphiperfol-5-ene, 7-β-(H)-silphiperfol-5-ene, α-campholene aldehyde, cameroonan-7-α-ol, silphiperfol-5-ene, silphiperfolan-7-β-ol, presilphiperfolan-9-α-ol and presilphiperfolan-8-ol (a major compound in *P. pallens*) (Table 2). Some of these compounds have previously only been recorded once in the essential oil literature, and have been isolated from *Eryophyllen staechadifolium* (König et al., 1997). Presilphiperfol-7-ene was first isolated from *Echinops giganteus* (Asteraceae) which is endemic to Cameroon and Nigeria (Weyerstahl et al., 1997a). It was also reported that cameroonan-7-α-ol also identified in *Pteronia* species contributed significantly to the strong woody amber-like odour of *E. giganteus*. α-Campholene aldehyde, another rare sesquiterpene (Fig. 1) was previously detected in the oil of *Juniperus oxycedrus* subsp. *macrocarpa* and *Salvia cyanescens* and represented 1.7–3.2% and 2.2% of the total oil composition, respectively (Sezik et al., 2005; Lucero et al., 2006). Presilphiperfolan-9-α-ol was reported in oils of *Artemisia chamaemelifolia* Vill. (Marco et al., 1996), *Artemisia laciniata* Willd. (Weyerstahl et al., 1997b) and *Haplopappus greenii* (Asteraceae) (Demirci et al., 2006), while presilphiperfolan-8-ol has been found in the hexane and ethanol rhizome extracts of *Silphium perfoliatum* and *Silphium integrifolium* in high levels (4.8–13.0%) (Kowalski, 2009).

In order to investigate variation between plants from the same population, three samples of *P. camphorata* were collected and the essential oil composition determined. The cluster analysis indicates that all three collections of *P. camphorata* are tightly grouped together (Fig. 2), with a very high correlation coefficient ($S_{\text{corr}} \approx 0.98$).

The variation between samples from different populations was also investigated for *P. pallens* which were collected at two different localities. The cluster analysis indicates that these samples all grouped together, with a high correlation ($S_{\text{corr}} \approx 1$) (Fig. 2), demonstrating that the oils are quantitatively and qualitatively similar in their composition for these species.

Table 2
Essential oil composition of the hydro-distilled essential oils as determined by GC-MS.

| RRI | Compound | <i>P. adenocarpa</i> | <i>P. camphorata</i> A | <i>P. camphorata</i> B | <i>P. camphorata</i> C | <i>P. elongata</i> | <i>P. empetrifolia</i> | <i>P. fasciculata</i> | <i>P. flexicaulis</i> | <i>P. glauca</i> | <i>P. glomerata</i> | <i>P. pallens</i> (Volmoed) | <i>P. pallens</i> (between Oudtshoorn and Volmoed) | <i>P. paniculata</i> | <i>P. viscosa</i> |
|------|--------------------------------|----------------------|---------------------------|---------------------------|---------------------------|--------------------|------------------------|-----------------------|-----------------------|------------------|---------------------|--------------------------------|---|----------------------|-------------------|
| | Yield | 0.43 | 0.29 | 0.19 | 0.71 | 0.15 | 0.48 | 0.14 | 0.11 | 0.51 | 0.2 | 0.28 | 0.55 | 0.34 | 0.28 |
| 952 | 1-Nonene | – | – | – | – | 0.5 | – | – | tr | – | – | – | – | – | – |
| 1032 | α-Pinene | 4.8 | 0.5 | 0.2 | 0.7 | 4.5 | 10.3 | 8.7 | 5.1 | 4.7 | 10.2 | 4.2 | 5.0 | 12.0 | 2.4 |
| 1035 | α-Thujene | – | 0.3 | – | 0.1 | – | – | – | – | – | 2.1 | – | – | – | – |
| 1048 | 2-Methyl-3-buten- 2-ol | – | – | – | – | 2.2 | – | – | – | – | – | – | – | – | – |
| 1118 | β-Pinene | 25.4 | 1.1 | 0.7 | 0.9 | 28.6 | 14.5 | 15.8 | 32.1 | 26.9 | 14.3 | 21.6 | 22.5 | 19.0 | 14.9 |
| 1132 | Sabinene | 3.3 | 9.1 | 7.1 | 12.7 | 7.8 | 2.5 | 2 | 2.8 | 33.6 | 13.3 | 32.2 | 30.8 | 22.0 | 9.0 |
| 1142 | Undecene | – | – | – | – | – | 0.4 | – | – | – | – | – | – | – | – |
| 1174 | Myrcene | 2.9 | – | – | – | 12.4 | 4.7 | 13.8 | 17.6 | 1.1 | 2.8 | 1.3 | 0.9 | 2.1 | 10.4 |
| 1176 | α-Phellandrene | – | 1.6 | 1.7 | 5.5 | – | – | – | – | – | – | – | – | – | – |
| 1188 | α-Terpinene | – | – | 0.1 | 0.3 | 0.7 | – | – | 0.5 | 0.7 | 1.3 | 0.3 | 0.6 | 0.7 | 0.7 |
| 1203 | Limonene | 2.8 | 5.0 | 3.8 | 7.7 | 3.0 | 11.2 | 2.7 | 6.1 | 3.3 | 3.2 | 4.6 | 4.3 | 13 | 2.7 |
| 1213 | 1,8-Cineole | 23.5 | 42.7 | 40.4 | 42.6 | – | 2.5 | – | – | – | 7.7 | – | – | – | 37 |
| 1218 | β-Phellandrene | – | – | – | – | – | 11.2 | – | 0.2 | 2.2 | – | 1.2 | 0.8 | 1.9 | 0.2 |
| 1246 | Z-(β)-Ocimene | 0.6 | 1.0 | 0.8 | 1.5 | 3.8 | 0.1 | – | – | – | 0.3 | 0.4 | – | – | – |
| 1255 | δ-Terpinene | 0.3 | 1.2 | 0.4 | 0.7 | – | – | – | – | – | – | – | 0.1 | – | 1.7 |
| 1255 | γ-Terpinene | – | – | – | – | 1.5 | – | – | 0.3 | – | 2.5 | – | – | 1.1 | – |
| 1266 | (E)-β-ocimene | 0.2 | 0.2 | tr | 0.1 | 1.4 | 0.3 | 0.9 | – | – | 1.1 | 1.0 | 1.4 | 0.5 | 0.9 |
| 1280 | p-Cymene | 14.3 | 17.1 | 21.1 | 10.0 | 3.6 | 9.8 | 9.2 | 1.3 | – | 9.9 | 0.5 | 1.1 | 2.9 | 1.7 |
| 1290 | Terpinolene | 0.2 | 0.3 | – | 0.1 | 0.5 | 0.1 | – | 0.3 | – | 0.6 | 0.1 | tr | 0.3 | 0.4 |
| 1327 | 3-Methyl-2-butenol | – | – | – | – | 0.6 | – | – | – | – | 0.1 | – | – | – | – |
| 1399 | Methyl-octanoate | – | – | – | – | 0.1 | 0.1 | – | – | – | 0.2 | – | – | – | 0.7 |
| 1406 | Presilphiperfol-7-ene | – | – | – | – | – | – | – | 0.4 | – | – | 0.4 | 0.3 | – | – |
| 1424 | 7-α-(H)- Silphiperfol-5-ene | – | – | – | – | – | – | – | – | – | – | – | 0.1 | – | – |
| 1435 | β-bourbonene | – | – | – | – | – | – | – | 0.5 | – | – | – | – | – | – |
| 1452 | 7 β-(H)- Silphiperfol-5-ene | – | – | – | – | – | – | – | 0.8 | – | – | – | 0.1 | – | – |
| 1466 | α-Cubebene | – | – | – | – | – | – | – | 0.5 | – | – | – | – | – | – |
| 1474 | trans-Sabinene hydrate | 0.1 | 0.4 | 0.8 | 0.6 | – | 0.1 | – | – | – | 0.4 | 1.1 | 1.4 | 1.2 | 0.2 |
| 1479 | Bicycloelemene | – | – | – | – | – | 0.1 | – | – | – | 0.1 | 1.1 | tr | 0.1 | – |
| 1495 | Silphiperfol-5-ene | – | – | – | – | – | – | – | 1.7 | – | – | 0.9 | 0.9 | – | – |
| 1499 | α-Campholene aldehyde | tr | – | – | – | – | 0.4 | – | – | – | – | – | tr | – | – |
| 1553 | Linalool | 0.1 | 0.4 | 2.8 | 3.1 | 0.2 | – | – | – | – | 0.1 | – | – | – | 0.1 |
| 1556 | cis-Sabinene hydrate | tr | – | – | – | – | – | – | – | – | 0.2 | 0.8 | 0.8 | 1.0 | – |
| 1571 | trans-p-Menth- 2-en-1-ol | 0.3 | 0.3 | 0.1 | 0.2 | 0.3 | 0.1 | – | tr | – | 0.4 | 0.1 | 0.1 | 0.5 | 0.3 |
| 1583 | Nopinone | 0.5 | – | – | – | 0.1 | 0.3 | – | – | – | 0.1 | 0.1 | 0.2 | – | – |
| 1586 | Pinocarvone | 0.7 | 0.1 | – | – | 0.1 | 0.6 | – | 0.1 | – | – | 0.1 | 0.4 | – | – |
| 1600 | Terpinen-4-ol | 3.6 | 5.8 | 3.1 | 2.4 | 7.0 | 0.8 | 6.3 | 2.3 | – | 9.5 | 2.2 | 2.4 | 9.2 | 8.4 |

Table 2 (continued)

| RRI | Compound | <i>P. adenocarpa</i> | <i>P. camphorata</i> A | <i>P. camphorata</i> B | <i>P. camphorata</i> C | <i>P. elongata</i> | <i>P. empetrifolia</i> | <i>P. fasciculata</i> | <i>P. flexicaulis</i> | <i>P. glauca</i> | <i>P. glomerata</i> | <i>P. pallens</i> (Volmoed) | <i>P. pallens</i> (between Oudtshoorn and Volmoed) | <i>P. paniculata</i> | <i>P. viscosa</i> |
|------|-------------------------------------|----------------------|---------------------------|---------------------------|---------------------------|--------------------|------------------------|-----------------------|-----------------------|------------------|---------------------|--------------------------------|---|----------------------|-------------------|
| | Yield | 0.43 | 0.29 | 0.19 | 0.71 | 0.15 | 0.48 | 0.14 | 0.11 | 0.51 | 0.2 | 0.28 | 0.55 | 0.34 | 0.28 |
| 1632 | <i>cis</i> -p-Menth-2-en-1-ol | 0.2 | – | – | – | 0.2 | 0.2 | – | tr | – | 0.3 | – | – | 0.3 | 0.2 |
| 1639 | <i>trans</i> -p-Menth-2,8-dien-1-ol | – | 0.4 | 0.4 | – | – | – | – | – | – | – | – | – | – | – |
| 1648 | Myrtenal | – | 0.1 | 0.2 | 0.1 | 0.3 | 0.8 | – | 0.1 | – | tr | 0.3 | 0.6 | – | – |
| 1651 | Sabinaketone | – | 0.3 | 0.2 | 0.2 | – | – | – | – | – | – | – | 0.6 | – | – |
| 1661 | <i>trans</i> -Pinocarveol | 2.3 | 0.2 | 0.2 | 0.1 | 0.4 | 1.5 | – | – | – | 0.1 | 0.5 | 1 | – | 0.1 |
| 1671 | Methyl chavicol (=estragol) | – | 4.5 | 6.8 | 2.9 | – | – | – | – | – | – | – | – | – | – |
| 1682 | α -Terpineol | 1.7 | 2.3 | 4.8 | 3.9 | 1.9 | 0.8 | 1.3 | 1.1 | – | 1.4 | 0.1 | 0.2 | 1.6 | 1.5 |
| 1687 | α -Humulene | – | – | – | – | – | – | – | – | – | – | 0.3 | 0.5 | – | – |
| 1689 | <i>trans</i> -Piperitol | – | – | – | – | – | – | – | – | – | 0.2 | – | – | – | 0.1 |
| 1690 | Cryptone | 0.4 | – | – | – | 0.2 | – | – | – | – | – | – | – | – | – |
| 1726 | Verbenone | 0.3 | – | – | – | – | 0.7 | – | – | – | – | – | – | – | – |
| 1726 | Germacrene D | – | – | – | – | – | – | – | 0.1 | – | – | 0.8 | 0.9 | 1.1 | – |
| 1740 | α -Muurolole | – | – | – | – | – | 1.2 | – | – | – | 0.1 | – | – | – | – |
| 1743 | α -Cadinene | 0.3 | – | – | – | – | – | – | 0.4 | – | – | – | – | – | – |
| 1751 | Carvone | 0.5 | 1.2 | 0.1 | 0.7 | – | – | – | – | – | – | – | – | – | – |
| 1755 | Bicyclogermacrene | – | – | – | – | 0.8 | 1.5 | – | 0.2 | – | 2.8 | 1.3 | 0.7 | 2.4 | – |
| 1773 | δ -Cadinene | 0.4 | 0.1 | – | 0.1 | – | 2.1 | – | 0.1 | – | 0.3 | tr | 0.1 | 0.8 | 0.3 |
| 1776 | γ -Cadinene | – | – | 0.1 | – | – | – | – | – | – | 0.1 | 0.1 | 0.2 | – | – |
| 1786 | Kessane | 0.1 | – | – | – | 0.1 | 1.5 | – | 0.1 | – | 0.5 | 0.6 | 0.7 | – | – |
| 1797 | p-Methyl acetophenone | 0.1 | 0.5 | 0.5 | – | – | – | – | – | – | – | – | – | – | – |
| 1802 | Cumin aldehyde | – | 0.4 | 0.4 | 0.2 | – | – | – | – | – | – | – | – | – | – |
| 1804 | Myrtenol | 1.4 | – | 0.2 | 0.1 | 0.3 | 0.8 | – | 0.2 | – | 0.2 | 0.4 | 0.7 | – | – |
| 1804 | Liguloxide | 0.3 | – | – | – | 0.1 | 1.4 | – | – | – | 0.2 | 0.5 | 0.6 | – | – |
| 1834 | <i>trans</i> -Carveol | 0.2 | 0.4 | 0.3 | 0.2 | – | 0.5 | – | – | – | – | 0.3 | 0.1 | – | – |

| | | | | | | | | | | | | | | |
|------|-------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1864 | p-cymen-8-ol | 0.8 | 0.2 | 0.4 | 0.2 | 0.1 | 0.2 | – | 0.1 | 0.1 | – | 0.1 | 0.1 | 0.1 |
| 1867 | Thymol acetate | – | – | – | – | – | 11.0 | – | – | – | – | – | – | – |
| 1916 | α -Agarofuran | – | – | – | – | – | 0.2 | – | 0.3 | – | 0.3 | 0.6 | – | – |
| 1924 | Silphiperfolan-7- β -ol | – | – | – | – | – | – | – | 0.1 | – | 0.1 | 0.1 | – | – |
| 1957 | Cubebene | – | – | – | – | – | – | – | 0.6 | – | – | – | – | – |
| 1976 | Cameroonan-7- α -ol | – | – | – | – | – | – | – | 0.6 | – | 0.4 | – | – | – |
| 2008 | Caryophyllene oxide | tr | 0.3 | 0.3 | 0.4 | 0.3 | 0.2 | 0.3 | 1.3 | – | 0.4 | 0.7 | – | – |
| 2030 | Methyl eugenol | – | – | 0.4 | 0.2 | – | – | – | – | – | – | – | – | – |
| 2030 | Methyl eugenol | 0.3 | – | – | – | – | – | – | – | – | – | – | – | – |
| 2024 | Presilphiperfolan-9- α -ol | – | – | – | – | – | – | – | 0.9 | – | – | – | – | – |
| 2037 | (E)-Nerolidol | – | – | – | – | – | – | 4 | 0.5 | – | – | tr | 0.2 | – |
| 2067 | Presilphiperfolan-8-ol | – | – | – | – | – | 0.9 | – | 0.9 | – | 7.5 | 7.9 | – | – |
| 2081 | Humulene epoxide III | 0.4 | – | tr | – | – | – | – | – | – | – | – | – | – |
| 2088 | 1-epi-Cubenol | – | – | – | – | 0.1 | 0.1 | – | 0.2 | – | – | – | – | – |
| 2098 | Globulol | – | – | – | – | 0.1 | – | – | – | 0.4 | – | tr | 0.1 | – |
| 2104 | Viridiflorol | 0.1 | – | – | – | 0.1 | 1.9 | 31.1 | 2.3 | 0.3 | – | – | 0.2 | – |
| 2113 | Cumin alcohol | 0.3 | 0.2 | 0.1 | 0.1 | 0.1 | – | – | – | – | 0.1 | 0.3 | – | – |
| 2127 | 10-Epi- γ -Eudesmol | tr | – | – | – | – | 0.7 | – | 5.4 | – | 8.8 | 6.4 | – | – |
| 2144 | Spathulenol | 2.4 | 0.1 | – | 0.1 | 3.6 | 4.7 | 2.1 | 1.3 | 8.6 | – | – | 2.3 | 2.2 |
| 2183 | Thymol | – | – | – | – | – | 1.7 | – | – | – | – | – | – | – |
| 2196 | T-muurolol | 0.3 | – | – | – | – | 0.9 | – | 0.4 | 0.2 | – | – | 0.2 | 0.1 |
| 2209 | Torreyol | – | – | – | – | – | 0.1 | – | – | 0.1 | – | – | – | – |
| 2232 | α -Bisabolol | – | – | – | – | – | 1.3 | – | – | – | – | – | – | – |
| 2247 | <i>trans</i> - α -Bergamotol | – | – | – | – | – | 0.1 | – | – | 0.6 | – | 0.1 | – | – |
| 2255 | α -Cadinol | 0.4 | 0.2 | – | – | – | 1.5 | 0.2 | – | 0.3 | 0.9 | 0.3 | – | 0.9 |
| 2264 | Intermedeol | – | – | – | – | – | – | – | – | 0.8 | – | – | – | – |
| | Total | 96.8 | 98.5 | 98.5 | 98.7 | 98.8 | 97.4 | 98.6 | 91.8 | 98.0 | 97.9 | 97.6 | 96.8 | 97.2 |

tr: trace amount ($\leq 0.05\%$).

RRI: relative retention index.




Fig. 1. Structure of rare sesquiterpenes identified in the oils of *Pteronia* species.




Fig. 2. Cluster analysis of essential oil composition.

Acknowledgements

Grant support by the National Research Foundation (South Africa) is acknowledged. The authors are also grateful to Mr. J. Vlok for the identification of plant species.

References

- Demirci, B., Baser, K.H.C., Tabanca, N., Wedge, D.E., 2006. Characterization of volatile constituents of *Haplopappus greenei* and studies on the antifungal activity against phytopathogens. *Journal of Agricultural and Food Chemistry* 54, 3146–3150.
- Goldblatt, P., Manning, J.C., 2000. *Cape Plants. A Conspectus of the Cape Flora of South Africa*. National Botanical Institute of South Africa and MBA Press, Missouri Botanical gardens, USA.
- König, W.A., Fricke, C., Melching, S., Wihstutz, S., 1997. Identification of rare sesquiterpene structures in essential oils. In: Başer, K.H.C., Kirimer, N. (Eds.), *Progress in Essential Oil Research*. InAnadolu University Press, Eskişehir.
- Kowalski, R., 2009. *Silphium* L. extracts — composition and protective effect on fatty acids content in sunflower oil subjected to heating and storage. *Food Chemistry* 112, 820–830.
- Lucero, M., Fredrickson, E.L., Estell, R.E., Morrison, A.A., Richman, D., 2006. Volatile composition of *Gutierrezia sarothrae* (broom snakeweed) as determined by steam distillation and solid phase microextraction. *Journal of Essential Oil Research* 18, 121–125.
- Marco, A.A., Sanz-Cervera, J.F., Morante, M.D., Garcia-Lliso, V., Valles-Xirau, J., Jakupovic, J., 1996. Tricyclic sesquiterpenes from *Artemisia chamaemelifolia*. *Phytochemistry* 41, 837–844.
- Pestana, M.H., Gageiro, J.N., 2000. *Análise de Dados para Ciências Sociais. A complementaridade do SPSS. Edições Sílabo, Lisboa*.
- Rohlf, J.F., 1992. *NTSYS-pc, Numerical Taxonomy and Multivariate Analysis System*. Applied Biostatistics Inc., New York.
- Sezik, E., Kocakulak, E., Başer, K.H.C., Özek, T., 2005. Composition of the essential oils of *Juniperus oxycedrus* subsp. *macrocarpa* from Turkey. *Chemistry of Natural Compounds* 41, 284–285.
- Shearing, D., 1997. *Karoo. South African Wild Flower Guide 6*. Botanical Society of South Africa, Kirstenbosch, Claremont.

- Van Wyk, B.-E., 2008. A review of Khoi-San and Cape Dutch medical ethnobotany. *Journal of Ethnopharmacology* 119, 331–341.
- Weyerstahl, P., Seelmann, I., Marschall, H., Menut, C., 1997a. Some novel tricyclic sesquiterpene skeletons from the essential oil of *Echinops giganteus*. In: Başer, K.H.C., Kirimer, N. (Eds.), *Proceedings of the 28th International Symposium on Essential Oils*. In Anadolu University Press, Eskisehir, pp. 166–181.
- Weyerstahl, P., Marschall, H., Schroder, M., Wahlburg, H.C., Kaul, V.K., 1997b. The sesquiterpene fraction of the essential of *Artemisia laciniata* Willd. *Flavour Fragrance Journal* 12, 315–325.
- Zedro, C., Jakupovic, J., Bohlmann, F., 1990. Diterpenes and other constituents from *Pteronia* species. *Phytochemistry* 29, 1231–1245.