

TUTDoR

Fictitious time integration method for solving the time fractional gas dynamics equation

Item Type	Article
Authors	Partohaghighi, Mohammad;Inc, Mustafa;Baleanu, Dimitru;Moshokoa, Seithuti Philemon
DOI	10.2298/TSCI190421365P
Publisher	National Library of Serbia
Rights	Attribution-NonCommercial-ShareAlike 4.0 International
Download date	2024-11-10 19:28:23
Item License	http://creativecommons.org/licenses/by-nc-sa/4.0/
Link to Item	https://hdl.handle.net/20.500.14519/726

FICTITIOUS TIME INTEGRATION METHOD FOR SOLVING THE TIME FRACTIONAL GAS DYNAMICS EQUATION

by

**Mohammad PARTOHAGHIGHI^a, Mustafa INC^{b*}, Dumitru BALEANU^c,
and Seithuti Philemon MOSHOKOA^d**

^a Department of Mathematics, University of Bonab, Bonab, Iran

^b Department of Mathematics, Science Faculty, Firat University, Elazig, Turkey

^c Department of Mathematics, Cankaya University, Balgat, Ankara, Turkey

^d Department of Mathematics and Statistics, Tshwane University of Technology,
Pretoria, South Africa

Original scientific paper

<https://doi.org/10.2298/TSCI190421365P>

In this work a powerful approach is presented to solve the time-fractional gas dynamics equation. In fact, we use a fictitious time variable y to convert the dependent variable $w(x, t)$ into a new one with one more dimension. Then by taking a initial guess and implementing the group preserving scheme we solve the problem. Finally four examples are solved to illustrate the power of the offered method.

Key words: *fictitious time integration method, group preserving scheme, time fractional gas dynamics equation, Caputo derivative*

Introduction

Calculus of fractional order is increasingly being worked to model various physical systems. Since many physical phenomena growing in engineering as well as in allied sciences can be depicted by developing models with the help of the fractional calculus. The fractional partial equations response ultimately converges to the non-fractional equations, fulfilling a notable care in the present times. The fractional derivatives are important due to broad scope of applications for mathematical modelling of problems such as traffic flow models, control, and relaxation processes [1-11]. There are some some analytical and numerical methods which are implemented to solve the fractional equations such as Group preserving scheme [12, 13], differential transform methods [14-16], homotopy perturbation methods [17-20]. This presented work is dedicated to study the following time fractional gas dynamics equation (TFGD):

$$\left\{ \begin{array}{l} {}^C D_{0^+}^\alpha w(x, t) + w(x, t)w_x(x, t) - w(x, t)(1 - w(x, t)) = \mathcal{K}(x, t) \\ w(x, 0) = g_1(x), x \in \Omega_x \\ w(x, T) = g_2(x), x \in \Omega_x \\ w(0, t) = h_1(t), t \in \Omega_t \\ w(b, t) = h_2(t), t \in \Omega_t \\ \Omega := \{(x, t) : a \leq x \leq b, 0 \leq t \leq T\} \end{array} \right. \quad (1)$$

* Corresponding author, e-mail: minc@firat.edu.tr

The gas dynamics equations are mathematical terminology that are adjunct on the physical laws of conservation such as the conservation of momentum, conservation of mass, and conservation of energy. Many authors solved the fractional gas dynamics equations using different numerical and analytical methods [21-29]. The differential transform method is implemented for solving TFGD [30, 31] and Fractional homotopy analysis transform method [32].

In this presented work, we create a powerful and reliable numerical approach to obtain the numerical solution of TFGD equation. This approach is firstly presented by Liu [33].

The fictitious time integration method (FTIM)

The Caputo fractional derivative of for fractional order $\alpha > 0$ is described by [34, 35]:

$${}^C \mathcal{D}_{0^+, t}^\alpha w(x, t) = \frac{\partial^\alpha w(x, t)}{\partial t^\alpha} = \begin{cases} \frac{1}{\Gamma(m-\alpha)} \int_0^t (t-\sigma)^{m-\alpha-1} \frac{\partial^m w(x, \sigma)}{\partial \sigma^m} d\sigma, & m-1 < \alpha < m \\ \frac{\partial w(x, t)}{\partial t^m}, & \alpha = m \end{cases} \quad (2)$$

By using Caputo fractional derivative definition and $0 \leq \alpha < 1$ for eq. (1):

$$\frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{w_\sigma(x, \sigma)}{(t-\sigma)^\alpha} d\sigma + ww_x - w(1-w) - \mathcal{K}(x, t) = 0 \quad (3)$$

Now, we multiply the eq. (3) into the parameter η as a fictitious damping coefficient which can help our to raise the stability of numerical integration:

$$\frac{\eta}{\Gamma(1-\alpha)} \int_0^t \frac{w_\sigma(x, \sigma)}{(t-\sigma)^\alpha} d\sigma + \eta ww_x - \eta w(1-w) - \eta \mathcal{K}(x, t) = 0 \quad (4)$$

Now, we impose the following transformation:

$$z(x, t, y) = (1+y)^d w(x, t), \quad 0 < d \leq 1 \quad (5)$$

By using previous transformation, eq. (4) converts to a new form:

$$\frac{\eta}{(1+y)^d} \left[\frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{z_\sigma(x, \sigma, y)}{(t-\sigma)^\alpha} d\sigma + \eta zz_x - \eta z(1-z) - \eta \mathcal{K}(x, t) \right] = 0 \quad (6)$$

From eq. (5) we can get:

$$\frac{\partial z}{\partial y} = d(1+y)^{d-1} w(x, t) \quad (7)$$

A combination of eqs. (7) and (6), concludes:

$$\frac{\partial z}{\partial y} = \frac{\eta}{(1+y)^d} \left[\frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{z_\sigma(x, \sigma, y)}{(t-\sigma)^\alpha} d\sigma + zz_x - z(1-z) \right] - \eta \mathcal{K}(x, t) + d(1+y)^{d-1} w \quad (8)$$

Then, eq. (8) can be transformed into a new form of PDE for z , by using $w = z / (1+y)^d$:

$$\frac{\partial z}{\partial y} = \frac{\eta}{(1+y)^d} \left[\frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{z_\sigma(x, \sigma, y)}{(t-\sigma)^\alpha} d\sigma + zz_x - z(1-z) \right] - \eta \mathcal{K}(x, t) + \frac{kz}{1+y} \quad (9)$$

By using:

$$\frac{\partial}{\partial y} \left[\frac{z}{(1+y)^d} \right] = \frac{z_y}{(1+y)^d} - \frac{dz}{(1+y)^{1+d}}$$

Next, by multiplying $1/(1+y)^d$ on both sides of eq. (9), we obtain:

$$\frac{\partial}{\partial y} \left(\frac{z}{(1+y)^d} \right) = \frac{\eta}{(1+y)^{2d}} \left[\frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{z_\sigma(x, \sigma, y)}{(t-\sigma)^\alpha} d\sigma + zz_x - z(1-z) \right] - \frac{\eta \mathcal{K}(x, t)}{(1+y)^d} \quad (11)$$

Using again the transformation $w = z/(1+y)^d$, we get:

$$w_y = \frac{\eta}{(1+y)^d} \left[\frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{z_s(x, \sigma, y)}{(t-\sigma)^\alpha} d\sigma + w(x, t, y)w_x(x, t, y) - w(x, t, y)[1 - w(x, t, y)] - \eta \mathcal{K}(x, t) \right] \quad (12)$$

We have to emphasize that y plays the fictitious co-ordinate role which able us to embed eq. (3) into a new PDE form in a space called 3-space, denoted R^3 . As well as, by a initially guess $w(x, t, 0)$, for all $y \geq 0$, $w = w(x, t, y)$ is an undetermined function with regard to the conditions in eq. (1).

Supposing $w_i^j(\zeta) := w(x_i, t_j, y)$ and $\mathcal{K}_i^j := \mathcal{K}(x_i, t_j)$ as the discrete values of w and \mathcal{K} at a point (x_i, t_j) . Implementing a semi-discretization to the eq. (12) concludes:

$$\frac{d}{dy} w_i^j(y) = \frac{\eta}{(1+y)^d} \cdot \left[\frac{1}{\Gamma(1-\alpha)} \int_0^{t_j} \frac{y_\sigma(x_i, \sigma, y)}{(t_j - \sigma)^\alpha} d\sigma + w_i^j(y) \frac{w_i^{j+1}(y) - w_i^j(y)}{\Delta x} - w_i^j(y)(1 - w_i^j(y)) - \mathcal{K}_i^j \right] \quad (13)$$

To calculate the aforementioned integral terms we can write the following approximation:

$$\int_0^{t_j} \frac{w_\sigma(x_i, \sigma, y)}{(t_j - \sigma)^\alpha} d\sigma \approx \sum_{l=1}^{j-1} \frac{w(x_i, t_{l+1}, y) - w(x_i, t_l, y)}{\Delta t (t_j - t_l)^\alpha} \quad (14)$$

Which stepsize Δx is $(b - a) / M_1$, $\Delta t = T / M_2$, $x_i = a + i\Delta x$ and $t_j = j\Delta t$.

The GPS for extracted system of ODE

In this stage, with $\mathbf{w} = (w_1^1, w_1^2, \dots, w_m^n)^T$ we can write the eq. (13) in the following form:

$$\mathbf{w}' = \mathbf{E}(\mathbf{w}, y), \quad \mathbf{w} \in \mathbb{R}^N, \quad y \in \mathbb{R} \quad (15)$$

where \mathbf{E} indicates a vector with ij -elements being the right-hand side of eq. (13) and \mathbf{w}' denotes the differential of \mathbf{w} with regard to y , and $N = M_1 \times M_2$ is the number of total grid point.

In this step we can use of group-preserving scheme (GPS) that introduced by Liu [33].

Let:

$$\mathbf{X}_{l+1} = \mathbf{B}(l)\mathbf{X}_l \quad (16)$$

where \mathbf{X}_l indicates the value of \mathbf{X} at the y_l and $\mathbf{B}(l)$ is a component of $SO_0(N, 1)$ which represents the group value of \mathbf{B} at y_l .

The Lie group can be created from \mathbf{C} which is a element of $so(N, 1)$:

$$\mathbf{B}_l = \exp[\Delta y \mathbf{C}(l)] = \begin{bmatrix} I_N + \frac{(\Psi_l - 1)}{\|\mathbf{E}_l\|^2} \mathbf{E}_l \mathbf{E}_l^T & \frac{\Phi_l \mathbf{E}_l}{\|\mathbf{E}_l\|} \\ \frac{\Phi_l \mathbf{E}_l^T}{\|\mathbf{E}_l\|} & \Psi_l \end{bmatrix}$$

where

$$\begin{aligned} \Psi_l &= \cosh\left(\frac{\Delta y \|\mathbf{E}_l\|}{\|\mathbf{w}_l\|}\right) \\ \Phi_l &= \sinh\left(\frac{\Delta y \|\mathbf{E}_l\|}{\|\mathbf{w}_l\|}\right) \end{aligned} \quad (17)$$

The $\mathbf{X} := (\mathbf{w}^T, \|\mathbf{w}\|)^T$ is a vector in Minkowskian space which converts eq. (15) into $\partial X / \partial y = \mathbf{C}\mathbf{X}$.

Where

$$\mathbf{C} := \begin{pmatrix} \mathbf{0}_{N \times N} & \frac{\mathbf{E}(\mathbf{w}, \theta)}{\|\mathbf{w}\|} \\ \frac{\mathbf{E}^T(\mathbf{w}, y)}{\|\mathbf{w}\|} & 0 \end{pmatrix} \in so(N, 1) \quad (18)$$

is a Liu algebra of the proper orthochronous Lorentz group $SO_0(N, 1)$. By replacing eq. (17) for B_l into eq. (16), we have:

$$\mathbf{w}_{l+1} = \mathbf{w}_l + \frac{(\Psi_l - 1)\mathbf{E}_l \mathbf{w}_l + \Phi_l \|\mathbf{w}_l\| \|\mathbf{E}_l\|}{\|\mathbf{E}_l\|^2} \mathbf{E}_l = \mathbf{w}_l + \Pi_l \mathbf{E}_l \quad (19)$$

in this stage, by selecting an initial value $u_i^j(0)$ we can apply GPS to solve numerical solution of eq. (15) from the initial fictitious y_0 to a chosen final fictitious time y_f . Moreover, we can control the convergence of w_i^j at the l and $l + 1$ steps by the following criterion:

$$\sqrt{\sum_{i,j=1}^{M_1, M_2} [\mathbf{w}_i^j(l+1) - \mathbf{w}_i^j(l)]^2} \leq \varepsilon \quad (20)$$

where ε is the convergence criterion.

Numerical examples

To show the power of our method four examples are solved.

Example 1: In order to show the ability of presented method we consider the following fractional TFGD equation with fractional order $\alpha = 0.9$.

$${}^C \mathcal{D}_{0^+}^\alpha w(x, t) + w(x, t) w_x(x, t) - w(x, t)(1 - w(x, t)) = 0$$

We implement the presented method to solve this problem under parameters $\eta = 35$ and $d = 0.1$. The initial guess and stepsize for y are supposed as $w_i^j(0) = 1e^{-5}$ and $\Delta y = 1e^{-3}$. We use the number of knots $M_1 = 25$ and $M_2 = 25$ in each co-ordinates of space and time, respectively. Also, considered domain in this example is $\Omega = [0, 1] \times [0, 1]$. Figure 1 is dedicated to show the exact solution $w(x, t) = e^{-x+t}$ and approximate solutions obtained by presented scheme. Power of the method with maximum absolute error $1.4 \cdot 10^{-17}$ is shown in fig. 2.

Example 2: Suppose following problem of TFGD with $\alpha = 1.5$ and $a = 2$.




Figure 1. Plots of the exact and approximate solutions for Example 1




Figure 2. Plot of error for Example 1

In order to manage the stability and convergency of the approach we choose $\eta=5, d=0.001$, respectively. Initial guess is $w_i^j(0) = 0.001$ and stepsize of method is same with Example 1. For $M_1 = M_2 = 39$ and $\Delta y = 10^{-10}$. Exact $w(x, t) = a^{t-x}$ and numerical solutions are plotted in fig. 3. Absolute numerical errors for this example $1 \cdot 10^{-17}$ which are depicted in fig. 4.

Example 3: We take the TFGD equation with:

$${}^C D_{0^+}^\alpha w(x, t) + w(x, t)w_x(x, t) - (1+t)^2 w^2(x, t) - x^2 = 0, \quad a > 0$$

Under parameters $\alpha = 0.3, \eta = 2, d = 0.001, \Delta y = 10^{-5}, M_1 = M_2 = 19$ and initial guess $w_i^j(0) = 0.0001$. The solutions and maximum absolute errors are demonstrated in figs. 5 and 6, respectively. Moreover, the exact solution of this example is $w(x, t) = x/(1+t)$ and $\Omega = [0, 1] \times [0, 1]$.




Figure 3. Plots of the exact and approximate solutions for Example 2




Figure 4. Plot of error for Example 2




Figure 6. Plot of error for Example 3





Figure 5. Plots of the exact and approximate solutions for Example 3

Conclusion

In this work we have converted TFGD equation into a new type of functional PDE in one more dimension by implementing a fictitious co-ordinate. Then by using a semi-discretization for original equation, the GPS as a geometric approach is imposed to solve the obtained system of first order ODE. Four numerical examples are solved, which demonstrate that our presented scheme is powerful and applicable to gain the numerical solutions of TFGD equation.

Nomenclature

B – an element of Lorentz group
 C – augmented matrix
 d – convergence rate parameter
 E – M -dimensional vector field in eq. (15)
 g_1 – initial solute concentration
 I_M – M -dimensional unit matrix
 h – boundary solute concentration
 \mathcal{K} – source term
 N – M number of discretized points
 $SO_0(M,1)$ – M -dimensional Lorentz group

T – time
 Δt – time stepsize
 w – solute concentration
 x – space dimension
 Δx – space stepsize

Greek symbols

α – fractional derivative order
 η – fictitious damping coefficient

References

- [1] Mainardi, A., Carpinteri, F., *Fractals and Fractional Calculus in Continuum Mechanics*, Springer Verlag, Wien, New York, 1997, pp. 277-290
- [2] Miller, K. S., B. Ross, B., *An Introduction the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, USA, 1993
- [3] Caputo, M., Mainardi, F., Linear Models of Dissipation in Anelastic Solids, *Rivista Del Nuovo Cimento*, 1 (1971), pp. 161-198
- [4] Oldham, K. B., Spanier, J., *The Fractional Calculus. Integrations and Differentiations of Arbitrary Order*, Academic Press, New York, USA, 1974
- [5] Samko, S. G., et al., *Fractional Integrals and Derivatives Theory and Applications*, Gordon and Breach, New York, USA, 1993
- [6] Podlubny, I., *Fractional Differential Equations*, Academic Press, San Diego, Cal., USA, 1999
- [7] Kilbas, A. A., et al., *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, The Netherlands, 2006
- [8] Hilfer, R., *Application of Fractional Calculus in Physics*, World Scientific, Singapore, 2000
- [9] Zaslavsky, G. M., *Hamiltonian Chaos and Fractional Dynamics*, Oxford University Press, Oxford, UK, 2005
- [10] Magin, R. L., *Fractional Calculus in Bioengineering*, Begell House Publisher, Inc., Danbury, Conn., USA, 2006
- [11] Gorenflo, R., Mainardi, F., Fractional calculus: Integral and Differential Equations of Fractional Order, in: *Fractals and Fractional Calculus*, SpringerVerlag, New York, USA, 1997
- [12] Hashemi, M. S., et al., A Lie Group Approach to Solve the Fractional Poisson Equation, *Rom. J. Phys.* 60, (2015), 9-10, pp. 1289-1297
- [13] Hashemi, M. S., et al., Solving the Time-Fractional Diffusion Equation Using a Lie Group Integrator, *Therma Science*, 19 (2015) Suppl. 1, pp. S77-S83
- [14] RaviKanth, A. S. V., Aruna, K., Differential Transform Method for Solving the Linear and Non-Linear Klein-Gordon Equation, *Comput. Phys. Commun.*, 180 (2009), 5, pp. 708-711
- [15] RaviKanth, A. S. V., Aruna, K., Differential Transform Method for Solving Linear and Non-Linear Systems of Partial Differential Equations, *Phys. Lett. A*, 372 (2008), 46, pp. 6896-6898
- [16] A.S.V. RaviKanth, K. Aruna, The 2-D Differential Transform Method for Solving Linear and Non-Linear Schro Dinger Equations, *Chaos, Solitons Fract.*, 41 (2009), 5, pp. 2277-2281
- [17] Kumar, S., Om P. S., Numerical Inversion of the Abel Integral Equation Using Homotopy Perturbation Method, *Z Naturforsch*, 65a, (2010), Sept., pp. 677-682
- [18] Kumar, S., H. et al., A Fractional Model of Gas Dynamics Equation and Its Analytical Approximate Solution by Using Laplace Transform, *Z Naturforsch*, 67, (2012), 1-2, pp. 389-396
- [19] Kumar, S., A Numerical Study for Solution of Time Fractional Non-Linear Shallow-Water Equation in Oceans, *Z Naturforsch A*, 68a, (2013), Sept., pp. 1-7
- [20] Kumar, S., Numerical Computation of Time-Fractional FokkerPlanck Equation Arising in Solid State Physics and Circuit Theory, *Z Naturforsch*, 68a (2013), Sept., pp. 1-8
- [21] Jafari, H., et al., Analytical Solution for Non-Linear Gas Dynamics Equation by Homotopy Analysis Method, *Appl. Appl. Math.*, 4 (2009), 1, pp. 149-154
- [22] Jawad, A. J. M. Applications of Hes Principles to Partial Differential Equations, *Appl. Math. Comput.*, 217 (2011), 16, pp. 7039-7047
- [23] Elizarova, T. G., *Quasi Gas Dynamics Equations*, Comput. Fluid Solid Mech., Springer-Verlag, Berlin, Germany, 2009
- [24] Evans, D. J., Bulut, H., A New Approach to the Gas Dynamics Equation: An Application of the Decomposition Method, *Int. J. Comput. Math.*, 79 (2002), 7, pp. 817-822
- [25] Steger, J. L., Warming, R. F., Flux Vector Splitting of the Inviscid Gas Dynamic Equations with Application Finite-Difference Methods, *Journal Comput. Phys.*, 40 (1981), 2, pp. 263-293
- [26] Aziz, A., Anderson, D., The Use of Pocket Computer in Gas Dynamics, *Comput. Educat.*, 9 (1985), 1, pp. 41-56
- [27] Rasulov, M., Karaguler, T., Finite Difference Scheme for Solving System Equation of Gas Dynamics in a Class of Discontinuous Function, *Appl. Math. Comput.*, 143 (2003), 1, pp. 145-164
- [28] Liu, T. P., *Non-Linear Waves in Mechanics and Gas Dynamics*, Defense Technical Information Center, Fort Belvoir, Va., USA, 1990, pp. 238-340
- [29] Tamsir, M., Srivastava, V. K., Revisiting the Approximate Analytical Solution of Fractional-Order Gas Dynamics Equation, *Alexandria Eng.*, 55 (2016), 2, pp. 867-74

- [30] Biazar, J., M. Eslami, M., Differential Transform Method for Non-Linear Fractional Gas Dynamics Equation, *Inter. J. Phys. Sci.*, 6 (2011), 5, pp. 12-03
- [31] Das, S., Kumar, R., Approximate Analytical Solutions of Fractional Gas Dynamics, *Appl. Math. Comput.*, 217 (2011), 24, pp. 9905-9915
- [32] Kumar, S., Rashidi, M. M., New Analytical Method for Gas Dynamics Equation Arising in Shock Fronts, *Comput. Phys. Commun.*, 185 (2014), 7, pp. 1947-1954
- [33] Liu, C.-S., Solving an Inverse Sturm-Liouville Problem by a Lie-Group Method, *Boundary Value Problems*, 2008 (2008), 1, pp. 749-865
- [34] Podlubny, I., *Fractional Differential Equations*, Academic Press, New York, USA, 1999
- [35] Debnath, L., Recent Applications of Fractional Calculus to Science and Engineering, *Int. J. Math Sci.*, 2003 (2003), 54, pp. 3413-3442